

Liquid
**Rotenone-
Pyrethrins**
Spray

C · o · n · c · e · n · t · r · a · t · e

ACTIVE INGREDIENTS:

Rotenone	1.10% W/W
Other Cube Extractives	1.10%
Pyrethrins	0.80%
OTHER INGREDIENTS	<u>97.00%</u>
TOTAL	100.00% W/W

Contains Aromatic Petroleum Solvent

KEEP OUT OF REACH OF CHILDREN

CAUTION

(See Back Panel for Additional Precautionary Statements and First Aid)

TRUSTED SINCE 1926

BONIDE®



A combination of insecticides for easy mixing with water. For use on the listed fruits, vegetables and ornamental plants. Also kills lice, fleas and ticks.

VERBKATE

LIQUID ROTENONE-PYRETHRINS SPRAY Concentrate

CONTROLS insect pests, such as aphids, leafhoppers, Japanese beetles, stink bugs, flea beetles...**USE** on apples, plums, peas, squash, cabbage, oak, elm, cedar, petunia, aster, geranium, marigold...**MIX** 1 to 4 tsp. per gallon of water. **Mixture ratios vary. Read and use according to label directions. APPLY** using a sprayer or watering can.

Store and transport in an upright position.

BN

EPA Est. No. 4-NY-1

EPA Reg. No. 4-315

Buyers guarantee limited to label claims

BONIDE PRODUCTS, INC.

6301 Suttiff Road
Oriskany, NY 13424

VERBKATE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Store and transport in an upright position. Store in a cool, dry area inaccessible to children or pets.

DISPOSAL: *If empty-* Do not reuse this container. Place in trash or offer for recycling if available. *If partially filled-* Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Page: One

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Mix this product with water as directed to control the listed pests of ornamentals, vegetables, fruit and berries. Use a hand or automatic sprayer as a coarse spray. Use sufficient spray mix to give complete coverage of the plant.

Page: Two

FRUITS AND BERRIES

APPLE, PEAR, PEACH, PLUM, CHERRY, GRAPE, CURRANT, GOOSEBERRY, RASPBERRY, BLACKBERRY, STRAWBERRY, AND BLUEBERRY.

For Aphids, Leafhoppers, Exposed Thrips mix 1 teaspoon with 1 gallon of water. For Imported Currant Worms, Gooseberry Fruit Worms, Raspberry Fruit Worms, Blueberry Maggots, Cherry Fruit Flies, Strawberry Root Worms (adult), Red-necked Borers, Japanese Beetles, and Mites mix 2 teaspoons with 1 gallon of water. For Strawberry Leafrollers and Rose Chafers mix 4 teaspoons with 1 gallon of water. Do not apply within one day of collection. Repeat every 5-10 days or as needed.

Page: Three

Page: Four

VEGETABLES

PEAS, BEAN, BEET, CUCUMBER, SQUASH, MUSKMELON, POTATO, TOMATO, CABBAGE, SPINACH, LETTUCE, CELERY, ASPARAGUS, PEPPER, EGGPLANT, ONION, CARROT AND CORN. For Aphids, Leafhoppers, and Exposed Thrips mix 1 teaspoon with 1 gallon of water. For Cabbage Worms, Bean Beetles, Beet Webworms, Sod Webworms, Asparagus Beetles, Japanese Beetles, Lygus Bugs, Whiteflies, Garden Fleahoppers, Harlequin Bugs, Obliquebanded Leaf Rollers, Stink Bugs, Celery Leaf-tiers, Melon Worms, Tortoise Beetles, Garden Webworms, Alfalfa Webworms and Mites mix 2 teaspoons with 1 gallon of water. For Colorado Potato Beetle, Cucumber Beetles, Flea Beetles, Vegetable Weevils, Squash Bugs (nymphs), Squash Vine Borers, European Corn Borers mix 4 teaspoons with 1 gallon of water. Do not apply within one day of harvest. Repeat every 5-10 days or as needed. European Corn Borers - Apply 3 to 4 applications or more as eggs start to hatch and repeated at 5 to 7 day intervals.

ORNAMENTALS

POPLAR, LINDEN, DOGWOOD, WILLOW, BIRCH, BOXELDER, MAPLE, OAK, ELM, PINE, SPRUCE, CEDAR, YEW, SPIREA, LILAC, ZINNIA, ASTER, PETUNIA, GERANIUM, DAHLIA, GLADIOLUS, PHLOX, SWEET PEA, MARIGOLD, CARNATION, CHRYSANTHEMUM AND ROSE. For Aphids, Leafhoppers, and Exposed Thrips mix 1 teaspoon with 1 gallon of water. For Oblique-banded Leaf Rollers, Sawfly larvae, Boxelder Bugs, Japanese Beetles, Elm Leaf Beetle larvae, Mites, Lygus Bugs, Lace Bugs, Greenhouse Whiteflies, Stink Bugs, Garden Fleahoppers, and Greenhouse Leaf-tiers mix 2 teaspoons with 1 gallon of water. For Cucumber Beetles, Rose Chafers, Ants, Aster Beetles, and Flea Beetles mix 4 teaspoons with 1 gallon of water. Do not apply within one day of collection. Repeat every 5-10 days or as needed.

Page: Five

Page: Six

FIRST AID

If swallowed	Immediately call a poison control center or doctor. Do not induce vomiting unless told to do by a poison control center or doctor. Do not give any liquid to the person. Do not give anything by mouth to an unconscious person.
If in eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If on skin or clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If inhaled	Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for treatment advice.

NOTE TO PHYSICIAN

May pose an aspiration pneumonia hazard. Contains petroleum distillate. Have the container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-858-7378 for emergency medical information.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed or inhaled. Causes moderate eye irritation. Avoid contact with eyes, clothing or breathing (spray mist). Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Do not apply directly to water. Runoff and drift from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.

Page: Seven

Page: Eight